



جامعة بورسعيد كلية التمريض



| Year 20 ⁷ 1-20 ⁷ 1 | Subject: palliative care |
|--|--------------------------|
| | First year |
| Date: / / 20 Y Y | Total Marks: ۲0 |
| Time allowed: | Dr. Hayat Mohamed |

"ALL Questions should be answered".

(75 Marks)

Part I

(8 Mark)

Read the following statement carefully, then circle (T) if statement is true and (F) if the statement is false.

| 1. | person who complain bypass diarrhea suspect constipation | (T) (F | 7) |
|----|--|---------|----|
| ۲ | A score of 5/12 indicates severe constipation and requires aggressive intervention to empty the colon. | (T) (F | 7) |
| ٣ | hypomagnesemia and hypocalcemia are causes diarrhea | T) (F) | |
| ٤ | Cachexia is defined as a loss of desire to eat or a loss of appetite associated with a decrease in food intake . | (T) (F) |) |
| ٥ | Nausea is a mechanism whereby the body tries to purge itself of noxious. | (T) (F | 7) |
| ٦ | Dexamethasone the drug of choice for mechanical and chemical causes of nausea | (T) (F | 7) |
| ٧ | Fatigue is always a sign of a serious medical condition. | (T) (F) |) |
| ٨ | Palliative care is best used for simple problems like colds and flus | (T) (F) |) |

Part II (VMarks)

Read the following statement carefully, and then choose one answer.

- 1. A client, 45 years old, has acute pain following a fractured ankle. The physician's order is acetaminophen (Tylenol) with 30 mg of codeine one to two tablets, q3—4h prn. Although the client had taken two tablets 1 hour ago, she reports having severe pain. What nursing action is most appropriate?
- a. Consult the nurse-in-charge
- **b.** Reassess the client's pain in 15 minutes
- c. Notify the physician of the client's pain level
- d. Administer an additional dose while awaiting a new order
- 2. A nurse is caring for a 50-year-old client who is clinically obese and presented to the emergency department with complaints of right-sided back pain. The nurse removes the bedpan and notes white, pasty stool. What would be the most appropriate action?
- a. Encouraging the client to increase daily fluid intake
- **b.** Encouraging the client to increase daily fibre intake
- c. Documenting the findings and notifying the health care provider
- **d.** Obtaining a thorough dietary history for the past 72 hours
- 3. Non-verbal communication all of the following except
- a. Reflect feeling
- **b.** Listening
- c. Talking
- **d.** Respect





جامعة بورسعيد كلية التمريض



4. Besides watery stools, which of these symptoms also may occur during diarrhea

- a. Abdominal cramping
- **b.** Fever
- **c.** Hypertension
- d. Chest pain

5. Step 2 (moderated pain) ladder analgesic according the WHO consist of

- a. Nonopioid adjuvants
- **b.** Weak opioid adjuvants
- c. Weak opioid, nonopioid adjuvants
- d. Strong opioid, nonopioid adjuvants

6. Neuralgic pain signs and symptoms is

- a. Constant pain that occasionally radiates
- **b.** Heat, burning, numbness or tingling over the area.
- c.Light pressure from non-painful stimuli can produce severe pain.
- d. Sudden attack of stabbing pain
- 7. The chief aim of palliative care is ------
- **a.** To cure illness
- **b.** To improve quality of life
- c. To hasten death
- d. To prolong life

Part IV (10 Marks)

Answer the following questions

- 1. Discuss the pharmacological management about dyspnea?
- 2. List the ten steps to breaking bad news?

Best wishes

Dr. Hayat Mohamed